

Critical and multicritical behavior of the $\pm J$ Ising model in two and three dimensions

M Hasenbusch¹, F Parisen Toldin^{2,3}, A Pelissetto⁴, E Vicari⁵

¹ Institut für Theoretische Physik, Universität Leipzig, D-04009 Leipzig, Germany

² Max-Planck-Institut für Metallforschung, D-70569 Stuttgart, Germany

³ Institut für Theoretische und Angewandte Physik, Universität Stuttgart, D-70569 Stuttgart, Germany

⁴ Dipartimento di Fisica dell'Università di Roma "La Sapienza" and INFN, I-00185 Roma, Italy

⁵ Dipartimento di Fisica dell'Università di Pisa and INFN, I-56127 Pisa, Italy

E-mail: parisen@mf.mpg.de

Abstract. We report our Monte Carlo results on the critical and multicritical behavior of the $\pm J$ Ising model [with a random-exchange probability $P(J_{xy}) = p\delta(J_{xy} - J) + (1-p)\delta(J_{xy} + J)$], in two and three dimensions. We study the transition line between the paramagnetic and ferromagnetic phase, which extends from $p = 1$ to a multicritical (Nishimori) point. By a finite-size scaling analysis, we provide strong numerical evidence that in three dimensions the critical behavior along this line belongs to the same universality class as that of the critical transition in the randomly dilute Ising model. In two dimensions we confirm that the critical behavior is controlled by the pure Ising fixed point and that disorder is marginally irrelevant, giving rise to universal logarithmic corrections. In both two and three dimensions, we also determine the location of the multicritical Nishimori point, as well as the renormalization-group dimensions of the operators that control the renormalization-group flow close to it.

1. Introduction

The $\pm J$ Ising model is a simplified model [1] for disordered uniaxial magnetic systems showing glassy behavior in some region of their phase diagram and represents an interesting theoretical laboratory to study the effects of quenched disorder and frustration. It is defined by the lattice Hamiltonian

$$\mathcal{H} = - \sum_{\langle xy \rangle} J_{xy} \sigma_x \sigma_y, \quad (1)$$

where $\sigma_x = \pm 1$, the sum is over the nearest-neighbor sites of a simple cubic lattice, and the exchange interactions J_{xy} are uncorrelated quenched random variables, taking values $\pm J$ with probability distribution

$$P(J_{xy}) = p\delta(J_{xy} - J) + (1-p)\delta(J_{xy} + J). \quad (2)$$

In the following we set $J = 1$ without loss of generality. For $p = 1$, we recover the standard Ising model, while for $p = 1/2$, we obtain the usual bimodal Ising spin-glass model. The phase diagram in two and three dimensions is sketched in Figure 1; it is symmetric for $p \rightarrow 1 - p$ and thus here and in the following we only consider the case $1 - p < 1/2$, i.e., $p > 1/2$.




Figure 1. Phase diagram of the $\pm J$ Ising model in the T - p plane, in dimensions $d = 2$ (left) and $d = 3$ (right).

In both $d = 2$ and $d = 3$, the model exhibits a paramagnetic-ferromagnetic (PF) phase transition in the region of low frustration. The PF transition line starts at the Ising point $X_{Is} = (T = T_{Is}, p = 1)$, where T_{Is} is the critical temperature of the Ising model, and extends up to the multicritical Nishimori point (MNP) at $X_{MNP} = (T^*, p^*)$. Along this line, the critical behavior is analogous to that observed in randomly dilute Ising (RDI) models.

In two dimensions, renormalization-group (RG) and conformal field theory predict [2, 3, 4, 5] that disorder is marginally irrelevant, thus the critical behavior is controlled by the pure Ising fixed point, with universal logarithmic corrections [6]. In three dimensions disorder is relevant and the model shows a second-order phase transition in the three-dimensional (3D) RDI universality class [7], which describes transitions in generic diluted Ising systems with ferromagnetic exchange interactions [8].

As argued in [9, 10, 11], the MNP is located along the so-called Nishimori line (N line) [12, 13] defined by the equation

$$\tanh \beta = 2p - 1, \quad (3)$$

where $\beta \equiv 1/T$. At the MNP the transition line is predicted to be parallel to the T axis [11]. Then, it reaches the $T = 0$ axis at $X_c = (0, p_c)$. It has been proved [12] that ferromagnetism can only exist in the region $p \geq p^*$; thus, p_c must satisfy the inequality $p_c \geq p^*$. Recent numerical works indicate that p_c is strictly larger than p^* , though deviations are quite small, both in two [14] and three [15] dimensions, as well as in related models [16, 17].

The critical behavior for $p < p^*$ depends on the dimension. The 3D $\pm J$ model exhibits a paramagnetic-glassy (PG) transition line, which extends from the MNP up to $p = 1/2$. On this transition line, the critical behavior is independent on p and belongs to the Ising spin-glass universality class [18]. In contrast, in two dimensions there is no evidence of a finite-temperature glassy phase. Glassy behavior is only expected for $T = 0$ and $p < p_c$: the glassy phase at $T = 0$ is unstable with respect to thermal fluctuations.

2. Finite-size scaling

2.1. Paramagnetic-ferromagnetic transition line

We present here the strategy used to analyze the critical behavior on the PF transition line in the 3D $\pm J$ model. The two-dimensional (2D) case is analyzed in [6].

According to the renormalization group (RG), in the case of periodic boundary conditions and for $L \rightarrow \infty$, where L is the lattice size, a generic RG invariant quantity R at the critical

temperature $1/\beta_c$ behaves as

$$R(L, \beta = \beta_c) = R^* (1 + c_{11}L^{-\omega} + c_{12}L^{-2\omega} + \dots + c_{21}L^{-\omega_2} + \dots), \quad (4)$$

where R^* is the universal infinite-volume limit and ω and ω_2 are the leading and next-to-leading correction-to-scaling exponents. In 3D RDI systems scaling corrections play an important role, since ω is quite small; indeed, we have $\omega = 0.29(2)$ and $\omega_2 = 0.82(8)$ [19, 8]. Instead of computing the various quantities at fixed Hamiltonian parameters, we keep a RG invariant quantity R fixed at a given value R_f [20]. This means that, for each L , we determine the pseudocritical inverse temperature $\beta_f(L)$ such that $R(\beta = \beta_f(L), L) = R_f$. All interesting thermodynamic quantities are then computed at $\beta = \beta_f(L)$. The pseudocritical inverse temperature $\beta_f(L)$ converges to β_c as $L \rightarrow \infty$. The value R_f can be specified at will, as long as R_f is taken between the high- and low-temperature fixed-point values of R . The choice $R_f = R^*$ (where R^* is the critical-point value) improves the convergence of β_f to β_c for $L \rightarrow \infty$; indeed $\beta_f - \beta_c = O(L^{-1/\nu})$ for generic values of R_f , while $\beta_f - \beta_c = O(L^{-1/\nu-\omega})$ for $R_f = R^*$.

We can then consider any other RG invariant quantity R_α at fixed $R = R_f$, i.e., $\bar{R}_\alpha(L) = R_\alpha(L, \beta_f(L))$. For $L \rightarrow \infty$, one can show that [8]

$$\bar{R}_\alpha(L) = \bar{R}^* (1 + b_{11}L^{-\omega} + b_{12}L^{-2\omega} + \dots + b_{21}L^{-\omega_2} + \dots), \quad (5)$$

$$\bar{R}'_\alpha(L) = a_0 L^{1/\nu} (1 + a_{11}L^{-\omega} + a_{12}L^{-2\omega} + \dots + a_{21}L^{-\omega_2} + \dots), \quad (6)$$

$$\bar{\chi}(L) \equiv \chi(L, \beta = \beta_f(L)) = d_0 L^{2-\eta} (1 + d_{11}L^{-\omega} + d_{12}L^{-2\omega} + \dots + d_{21}L^{-\omega_2} + \dots) + d_b, \quad (7)$$

where $\bar{R}'_\alpha$ is the derivative of $\bar{R}_\alpha$ with respect to β and $\bar{\chi}$ is the susceptibility at fixed $R = R_f$. More details on this method can be found in [20, 8].

2.2. Multicritical Nishimori point

In the absence of external fields, the critical behavior at the MNP is characterized by two relevant RG operators. The singular part of the disorder-averaged free energy in a volume L^d can be written as

$$F_{\text{sing}}(T, p, L) = L^{-d} f(u_1 L^{y_1}, u_2 L^{y_2}, \{u_i L^{y_i}\}), \quad i \geq 3, \quad (8)$$

where $y_1 > y_2 > 0$, $y_i < 0$ for $i \geq 3$, u_i are the corresponding scaling fields, $u_1 = u_2 = 0$ at the MNP. The scaling fields u_i are analytic functions of the model parameters T and p . Using symmetry arguments, [10, 11] showed that the scaling axis corresponding to $u_2 = 0$ is along the N line, so that $u_2 = \tanh\beta - 2p + 1$. As for the scaling axis $u_1 = 0$, $\epsilon \equiv 6 - d$ expansion calculations predict it [11] to be parallel to the T axis. The extension of this result to lower dimensions suggests $u_1 = p - p^*$. Note that, if this conjecture holds, only the scaling field u_2 depends on the temperature T .

These results give rise to the following predictions for the FSS behavior around T^*, p^* . Along the N line, an RG invariant quantity R has the following behavior for $L \rightarrow \infty$:

$$R_N = R^* + b_{11}u_1 L^{y_1} + \dots, \quad (9)$$

where the subscript N indicates that R is restricted to the N line and we have neglected scaling corrections. Its derivative R' with respect to β behaves as

$$R' = b_{11}u'_1 L^{y_1} + b_{21}u'_2 L^{y_2} + \dots = b_{21}u'_2 L^{y_2} + \dots, \quad (10)$$

where the last equivalence holds only if u_1 does not depend on the temperature. This result gives us a method to verify the conjecture of [11]: once y_1 has been determined from the scaling behavior of a RG invariant quantity R close to the MNP, it is enough to check the scaling behavior of R' . If R' scales as L^ζ with $\zeta < y_1$, the conjecture is confirmed and $y_2 = \zeta$. Finally, along the N line the susceptibility χ is expected to behave as

$$\chi_N = d_0 L^{2-\eta} (1 + d_{11}u_1 L^{y_1} + \dots). \quad (11)$$

3. Results

3.1. The 3D $\pm J$ model at the paramagnetic-to-ferromagnetic transition line

In [7] we perform MC simulations of Hamiltonian (1) at $d = 3$ close to the PF transition line. Using the method outlined in Sec. 2.1, we fit the large- L limit of various RG-invariant quantities at fixed $R_\xi \equiv \xi/L = 0.5943$, where ξ is the correlation length; this value is very close to the fixed-point value $R_\xi^* = 0.5944(7)$ [8] at β_c . Our FSS analysis provides strong evidence that the critical behavior of the 3D $\pm J$ Ising model along the PF line belongs to the 3D RDI universality class. Indeed, all the RG-invariant quantities considered are in agreement with the results reported in [8]. Moreover, we find $\nu = 0.682(3)$ and $\eta = 0.036(2)$, in good agreement with the presently most accurate estimates [8] $\nu = 0.683(2)$ and $\eta = 0.036(1)$ for the 3D RDI universality class.

We also note that the random-exchange interaction in the $\pm J$ Ising model gives rise to frustration, while the RDI universality class describes transitions in generic diluted Ising systems with ferromagnetic exchange interactions. Therefore, our results imply that frustration is irrelevant along the PF transition line. Moreover, the observed scaling corrections are consistent with the RDI correction-to-scaling exponents $\omega = 0.29(2)$ and $\omega_2 = 0.82(8)$ [19, 8]. Thus, frustration does not introduce new irrelevant perturbations with RG dimension $|y_f| \lesssim 1$.

3.2. Multicritical Nishimori point

In [14, 15], we perform MC simulations of Hamiltonian (1) at $d = 2, 3$, along the N line defined by (3). In both cases, the position of the MNP as well as the RG dimension y_1 of the leading relevant operator u_1 are determined by fitting several RG-invariant quantities R to (9) (in the 2D case we also consider scaling corrections). We verify the conjecture that u_1 does not depend on T and compute the RG dimension associated with u_1 and u_2 .

For the 2D model, the MNP is located at $p^* = 0.89081(7)$, $T^* = 0.9528(4)$, the RG dimensions of the operators that control the multicritical behavior are $y_1 = 0.655(15)$ and $y_2 = 0.250(2)$, and the susceptibility exponent is $\eta = 0.180(5)$. In three dimensions, we find $p^* = 0.76820(4)$, $T^* = 1.6692(3)$, $y_1 = 1.02(5)$, $y_2 = 0.61(2)$, $\eta = -0.114(3)$.

References

- [1] Edwards S F, Anderson P W 1975 *J. Phys. F: Met. Phys.* **5** 965
- [2] Shalaev B N 1984 *Sov. Phys. Solid State* **26** 1811
- [3] Shankar R 1987 *Phys. Rev. Lett.* **58** 2466
Shankar R 1987 *Phys. Rev. Lett.* **59** 380
Ludwig A W W 1988 *Phys. Rev. Lett.* **61** 2388
Ceccatto H A, Naon C 1988 *Phys. Rev. Lett.* **61** 2389
Ludwig A W W 1989 *Phys. Rev. Lett.* **62** 980
- [4] Ludwig A W W 1990 *Nucl. Phys. B* **330**
- [5] Shalaev 1994 *Phys. Rep.* **237** 129
- [6] Hasenbusch M, Parisen Toldin F, Pelissetto A, Vicari E 2008 *Phys. Rev. E* **78** 011110
- [7] Hasenbusch M, Parisen Toldin F, Pelissetto A, Vicari E 2007 *Phys. Rev. B* **76** 094402
- [8] Hasenbusch M, Parisen Toldin F, Pelissetto A, Vicari E 2007 *J. Stat. Mech.: Theory Expt.* P02016
- [9] Georges A, Hansel D, Le Doussal P, and Bouchaud J 1985 *J. Phys. (Paris)* **46** 1827
- [10] Le Doussal P and Harris A B 1988 *Phys. Rev. Lett.* **61** 625
- [11] Le Doussal P and Harris A B 1989 *Phys. Rev. B* **40** 9249
- [12] Nishimori H 1981 *Prog. Theor. Phys.* **66** 1169
- [13] Nishimori H 2001 *Statistical Physics of Spin Glasses and Information Processing: An Introduction* (Oxford: Oxford University Press)
- [14] Hasenbusch M, Parisen Toldin F, Pelissetto A, Vicari E 2008 *Phys. Rev. E* **77** 051115
- [15] Hasenbusch M, Parisen Toldin F, Pelissetto A, Vicari E 2007 *Phys. Rev. B* **76** 184202
- [16] Wang C, Harrington J, Preskill J 2003 *Ann. Phys.* **303** 31
- [17] Ohzeki M, Nishimori H, Nihat Berker A 2008 *Phys. Rev. E* **77** 061116
- [18] Hasenbusch M, Pelissetto A, Vicari E 2008 *J. Stat. Mech.: Theory Expt.* L02001
- [19] Hasenbusch M, Pelissetto A, Vicari E 2007 *J. Stat. Mech.: Theory Expt.* P11009
- [20] Hasenbusch M 1999 *J. Phys. A: Math. Gen.* **32** 4851